



**Material suplementario**

---

**Comparación entre tres sistemas de puntuación de hemorragia en pacientes con fibrilación auricular no valvular. ¿Qué puede ofrecer la nueva puntuación ORBIT?**

---

**Supplementary Table 1**

## Bleeding Risk Stratification With ORBIT Score

<b>ORBIT risk score</b>	<b>Score</b>
<b>O</b> lder than 74 y	1
<b>R</b> educed hemoglobin [less than 13 g/dL and 12 g/dL in men and women, respectively]	2
<b>B</b> leeding history	2
<b>I</b> nsufficient kidney function [estimated glomerular filtration rate < 60 mL/min/1.73 m <sup>2</sup> ]	1
<b>T</b> reatment with antiplatelets	1
Maximum score	7

**Supplementary Table 2**

## Bleeding Risk Stratification With HAS-BLED and ATRIA

<b>HAS-BLED risk factor</b>	<b>Score</b>	<b>ATRIA bleeding risk factor</b>	<b>Score</b>
Hypertension (systolic blood pressure > 160mm Hg)	1	Anemia (ie, hemoglobin < 13 g/dL in men and < 12 g/dL in women and/or thrombocytopenia)	3
Abnormal renal and/or liver function	1 or 2	Severe renal disease (estimated glomerular filtration rate < 30 mL/min or dialysis dependent)	3
Stroke	1	Age ≥75 y	2
Bleeding tendency or predisposition	1	Prior hemorrhage	1
Labile international normalized ratios	1	Hypertension	1
Elderly (eg, age > 65 y, frail condition)	1		
Drugs (eg, concomitant antiplatelet or Nonsteroidal anti-inflammatory drugs) or alcohol excess/abuse	1 or 2		
Maximum score	9		10