

SUPPLEMENTARY MATERIAL

Table 1 of the supplementary material

Recalculated Predictive Model After Exclusion of Patients Who Died During Hospital Admission

| | OR (95%CI) | <i>P</i> |
|---|--------------------|----------|
| non-Shockable rhythm | 11.79 (3.16-18.63) | <.01 |
| Age (one additional year) | 1.05 (1.01-1.12) | .01 |
| Lactate level (one additional mmol/L) | 1.23 (1.09-1.43) | .04 |
| Time Elapsed to return of spontaneous circulation | 1.02 (1.00-1.06) | .03 |
| Diabetes | 3.65 (1.56-12.54) | .02 |

95%CI, confidence interval; OR, adjusted odds ratio.

Table 2 of the supplementary material


Recalculated Predictive Model After Exclusion of Patients With Noncardiac Causes of out-of-Hospital Cardiac Arrest

| | OR (95%CI) | <i>P</i> |
|---|---------------------|----------|
| non-Shockable rhythm | 43.01 (39.43-47.08) | <.01 |
| Age (1 additional year) | 1.05 (1.01-1.09) | .03 |
| Lactate level (one additional mmol/L) | 1.55 (1.20-2.00) | <.01 |
| Time Elapsed to return of spontaneous circulation | 1.05 (1.00-1.09) | .04 |
| Diabetes | 6.57 (1.55-12.93) | .01 |

95%CI, confidence interval; OR, adjusted odds ratio.

Figure of the supplementary material

Receiver operator characteristic curve for predictive model applied to in hospital survivors (n = 115).



95%CI, confidence interval; ROC, receiver operator characteristic.