

SUPPLEMENTARY DATA

Table 1

Association Between 1-year Posttransplant Serum Galectin-3 Levels (Continuous Variable) and the Risk of Death or Graft Failure over Long-Term Follow-up: Univariable and Multivariable Cox Regression Analyses

	Univariable analysis		Multivariable analysis	
	HR	95%CI	HR	95%CI
Recipient age, y	0.99	0.96–1.02	0.96	0.93–1
Donor age, y	1	0.97–1.02	–	–
Male recipient	0.85	0.39–1.86	–	–
Male donor	0.50	0.23–1.07	0.33	0.14–0.75
Right atrial pressure, mmHg	1.12	1.04–1.20	1.13	1.05–1.22
Left ventricular ejection fraction, %	0.97	0.92–1.02	–	–
Glomerular filtration rate, mL/min/m ²	1	0.98–1.02	–	–
Use of tacrolimus	1.94	0.88–4.24	2.87	1.18–6.98
Use of mycophenolate mophetil	1.13	0.43–2.96	–	–
Serum galectin-3 (ng/mL)	1.04	1.01–1.06	1.04	1.01–1.08

95%CI, 95%confidence interval; HR, hazard ratio.

Table 2

Association between 1-Year Posttransplant Serum Galectin-3 Levels ≥ 17.8 ng/mL (Dichotomous Variable) and the Risk of Death or Graft Failure Over Long-Term Follow-up: Univariable and Multivariable Cox Regression Analyses

	Univariable analysis		Multivariable analysis	
	HR	95%CI	HR	95%CI
Recipient age, y	0.99	0.96–1.02	0.96	0.93–1
Donor age, y	1	0.97–1.02	–	–
Male recipient	0.85	0.39–1.86	–	–
Male donor	0.50	0.23–1.07	0.35	0.15–0.80
Right atrial pressure, mmHg	1.12	1.04–1.20	1.10	1.02–1.19
Left ventricular ejection fraction, %	0.97	0.92–1.02	–	–
Glomerular filtration rate, mL/min/m ²	1	0.98–1.02	–	–
Use of tacrolimus	1.94	0.88–4.24	2.64	1.07–6.52
Use of mycophenolate mophetil	1.13	0.43–2.96	–	–
Serum galectin-3 ≥ 17.8 , ng/mL	2.96	1.44–6.10	2.74	1.07–7.02

95%CI, 95%confidence interval; HR, hazard ratio.